


Algorithms for graph visualization


Layouts for planar graphs. Shift method.

WINTER SEMESTER 2013/2014



Tamara Mchedlidze – MARTIN NÖLLENBURG





- Straight line drawing of a planar graph



- Straight line drawing of a planar graph





- Straight line drawing of a planar graph



Theorem [Wagner '36, Fary '48, Stein '51]

Every planar graph has a planar straight-line drawing.

- Straight line drawing of a planar graph



Theorem [Wagner '36, Fary '48, Stein '51]

Every planar graph has a planar straight-line drawing.

- These algorithms produce drawings with area **not bounded** by any polynomial on n .

This lecture:

Theorem [De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of a size $(2n - 4) \times (n - 2)$.

Next lecture:

Theorem [Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of a size $(n - 2) \times (n - 2)$.

Definition: Canonical Ordering

Let $G = (V, E)$ be a triangulated planar embedded graph of $n \geq 3$ vertices. An ordering $\pi = (v_1, v_2, \dots, v_n)$ is called a **canonical ordering**, if the following conditions hold for each k , $3 \leq k \leq n$.

- (C1) Vertices $\{v_1, \dots, v_k\}$ induce a 2-connected internally triangulated graph, call it G_k

Definition: Canonical Ordering

Let $G = (V, E)$ be a triangulated planar embedded graph of $n \geq 3$ vertices. An ordering $\pi = (v_1, v_2, \dots, v_n)$ is called a **canonical ordering**, if the following conditions hold for each k , $3 \leq k \leq n$.


- (C1) Vertices $\{v_1, \dots, v_k\}$ induce a 2-connected internally triangulated graph, call it G_k
- (C2) Edge (v_1, v_2) belongs to the outer face of G_k

Definition: Canonical Ordering


Let $G = (V, E)$ be a triangulated planar embedded graph of $n \geq 3$ vertices. An ordering $\pi = (v_1, v_2, \dots, v_n)$ is called a **canonical ordering**, if the following conditions hold for each k , $3 \leq k \leq n$.

- (C1) Vertices $\{v_1, \dots, v_k\}$ induce a 2-connected internally triangulated graph, call it G_k
- (C2) Edge (v_1, v_2) belongs to the outer face of G_k
- (C3) If $k < n$ then vertex v_{k+1} lies in the outer face of G_k , and all neighbors of v_{k+1} in G_k appear on the boundary of G_k consecutively.


Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering




Example of Canonical Ordering



Example of Canonical Ordering



Example of Canonical Ordering



Lemma

Every triangulated plane graph has a canonical ordering.

- Let $G_n = G$, and let v_1, v_2, v_n be the vertices of the outer face of G_n . Conditions C1-C3 hold.

Lemma


Every triangulated plane graph has a canonical ordering.

- Let $G_n = G$, and let v_1, v_2, v_n be the vertices of the outer face of G_n . Conditions C1-C3 hold.
- Induction hypothesis: vertices v_{n-1}, \dots, v_{k+1} have been chosen such that conditions C1-C3 hold for $k + 1 \leq i \leq n$.

Lemma

Every triangulated plane graph has a canonical ordering.


- Let $G_n = G$, and let v_1, v_2, v_n be the vertices of the outer face of G_n . Conditions C1-C3 hold.
- Induction hypothesis: vertices v_{n-1}, \dots, v_{k+1} have been chosen such that conditions C1-C3 hold for $k+1 \leq i \leq n$.
- Consider G_k . We search for v_k .



Lemma

Every triangulated plane graph has a canonical ordering.


- Let $G_n = G$, and let v_1, v_2, v_n be the vertices of the outer face of G_n . Conditions C1-C3 hold.
- Induction hypothesis: vertices v_{n-1}, \dots, v_{k+1} have been chosen such that conditions C1-C3 hold for $k+1 \leq i \leq n$.
- Consider G_k . We search for v_k .



Lemma

Every triangulated plane graph has a canonical ordering.

- Let $G_n = G$, and let v_1, v_2, v_n be the vertices of the outer face of G_n . Conditions C1-C3 hold.
- Induction hypothesis: vertices v_{n-1}, \dots, v_{k+1} have been chosen such that conditions C1-C3 hold for $k+1 \leq i \leq n$.
- Consider G_k . We search for v_k .




v_k should not be adjacent to a chord

Lemma

Every triangulated plane graph has a canonical ordering.

- Let $G_n = G$, and let v_1, v_2, v_n be the vertices of the outer face of G_n . Conditions C1-C3 hold.
- Induction hypothesis: vertices v_{n-1}, \dots, v_{k+1} have been chosen such that conditions C1-C3 hold for $k+1 \leq i \leq n$.
- Consider G_k . We search for v_k .




v_k should not be adjacent to a chord

Is it sufficient?


Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

Statement If v_k is not adjacent to a chord then removal of v_k leaves the graph biconnected.




Canonical Ordering Existence

- Why a vertex not adjacent to a chord exists?



Canonical Ordering Existence

- Why a vertex not adjacent to a chord exists?



Algorithm CO

```
forall the  $v \in V$  do  
   $\lfloor$  chords( $v$ )  $\leftarrow$  0; out( $v$ )  $\leftarrow$  false; mark( $v$ )  $\leftarrow$  false;  
  out( $v_1$ ), out( $v_2$ ), out( $v_n$ )  $\leftarrow$  true;  
for  $k = n$  to 3 do  
  choose  $v \neq v_1, v_2$  such that mark( $v$ ) = false, out( $v$ ) = true,  
    chords( $v$ ) = 0;  
   $v_k \leftarrow v$ ; mark( $v$ )  $\leftarrow$  true;  
  // Let  $w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2$  denote the boundary of  $G_{k-1}$ ;  
  // and let  $w_p, \dots, w_q$  be the unmarked neighbors  $v_k$ ;  
  out( $w_i$ )  $\leftarrow$  true for all  $p < i < q$ ;  
  update number of chords for  $w_i$  and its neighbors;
```

- chord(v) - number of chords adjacent to v
- mark(v) = true iff vertex v was numbered
- out(v)=true iff v is the outer vertex of current plane graph


Algorithm CO

```
forall the  $v \in V$  do  
   $\lfloor$  chords( $v$ )  $\leftarrow$  0; out( $v$ )  $\leftarrow$  false; mark( $v$ )  $\leftarrow$  false;  
  out( $v_1$ ), out( $v_2$ ), out( $v_n$ )  $\leftarrow$  true;  
for  $k = n$  to 3 do  
  choose  $v \neq v_1, v_2$  such that mark( $v$ ) = false, out( $v$ ) = true,  
    chords( $v$ ) = 0;  
   $v_k \leftarrow v$ ; mark( $v$ )  $\leftarrow$  true;  
  // Let  $w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2$  denote the boundary of  $G_{k-1}$ ;  
  // and let  $w_p, \dots, w_q$  be the unmarked neighbors  $v_k$ ;  
  out( $w_i$ )  $\leftarrow$  true for all  $p < i < q$ ;  
  update number of chords for  $w_i$  and its neighbors;
```


Lemma

Algorithm CO computes a canonical ordering of a graph in $O(n)$ time.

De Fraysseix Pach Pollack (Shift) Algorithm




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm




Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1





De Fraysseix Pach Pollack (Shift) Algorithm



Algorithm constraints: G_{k-1} is drawn such that


- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1






Algorithm constraints: G_{k-1} is drawn such that

- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1




De Fraysseix Pach Pollack (Shift) Algorithm





Algorithm constraints: G_{k-1} is drawn such that




- v_1 is on $(0, 0)$, v_2 is on $(2k - 6, 0)$
- Boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn x -monotone
- Each edge of the boundary of G_{k-1} (minus edge (v_1, v_2)) is drawn with slopes ± 1






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm







De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm




De Fraysseix Pach Pollack (Shift) Algorithm







De Fraysseix Pach Pollack (Shift) Algorithm







$L(10)$







De Fraysseix Pach Pollack (Shift) Algorithm







De Fraysseix Pach Pollack (Shift) Algorithm







De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm







De Fraysseix Pach Pollack (Shift) Algorithm






De Fraysseix Pach Pollack (Shift) Algorithm




De Fraysseix Pach Pollack (Shift) Algorithm




De Fraysseix Pach Pollack (Shift) Algorithm




De Fraysseix Pach Pollack (Shift) Algorithm




De Fraysseix Pach Pollack (Shift) Algorithm






- Each internal vertex is covered exactly once
- Coverance relation defines a tree in G
- But a forest in $G_i, 1 \leq i \leq n-1$




- Each internal vertex is covered exactly once
- Coverance relation defines a tree in G
- But a forest in $G_i, 1 \leq i \leq n-1$




- Each internal vertex is covered exactly once
- Coverage relation defines a tree in G
- But a forest in $G_i, 1 \leq i \leq n-1$

De Fraysseix Pach Pollack (Shift) Algorithm




- Each internal vertex is covered exactly once
- Coverance relation defines a tree in G
- But a forest in $G_i, 1 \leq i \leq n-1$

De Fraysseix Pach Pollack (Shift) Algorithm



- Each internal vertex is covered exactly once
- Coverance relation defines a tree in G
- But a forest in $G_i, 1 \leq i \leq n-1$



- Each internal vertex is covered exactly once
- Coverance relation defines a tree in G
- But a forest in $G_i, 1 \leq i \leq n-1$

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .
- Assume that this is true for G_{k-1} .

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .
- Assume that this is true for G_{k-1} .
- Let $w_1, \dots, w_p, v_k, w_q, \dots, w_t$ be the boundary of G_k .

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .
- Assume that this is true for G_{k-1} .
- Let $w_1, \dots, w_p, v_k, w_q, \dots, w_t$ be the boundary of G_k .
- Let $\delta_1 \leq \dots \leq \delta_p \leq \delta \leq \delta_q \leq \dots \leq \delta_t$.

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .
- Assume that this is true for G_{k-1} .
- Let $w_1, \dots, w_p, v_k, w_q, \dots, w_t$ be the boundary of G_k .
- Let $\delta_1 \leq \dots \leq \delta_p \leq \delta \leq \delta_q \leq \dots \leq \delta_t$.
- We set $\delta'_i = \delta_i$ for $1 \leq i \leq p$,
- $\delta'_i = \delta$ for $p + 1 \leq i \leq q - 1$ (for the neighbors of v_k)
- $\delta'_i = \delta_i$ for $q \leq i \leq t$.

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .
- Assume that this is true for G_{k-1} .
- Let $w_1, \dots, w_p, v_k, w_q, \dots, w_t$ be the boundary of G_k .
- Let $\delta_1 \leq \dots \leq \delta_p \leq \delta \leq \delta_q \leq \dots \leq \delta_t$.
- We set $\delta'_i = \delta_i$ for $1 \leq i \leq p$,
- $\delta'_i = \delta$ for $p + 1 \leq i \leq q - 1$ (for the neighbors of v_k)
- $\delta'_i = \delta_i$ for $q \leq i \leq t$.
- By induction hypothesis we can move $w_1 \dots, w_t$ by $\delta'_1 \dots \delta'_t$, respectively.

Lemma

Let $0 < \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$. If we shift $L(w_i)$ by δ_i to the right, we get a planar straight line drawing.

Proof

- The proof is by induction on i , i.e. we consider G_3, \dots, G_n .
- Assume that this is true for G_{k-1} .
- Let $w_1, \dots, w_p, v_k, w_q, \dots, w_t$ be the boundary of G_k .
- Let $\delta_1 \leq \dots \leq \delta_p \leq \delta \leq \delta_q \leq \dots \leq \delta_t$.
- We set $\delta'_i = \delta_i$ for $1 \leq i \leq p$,
- $\delta'_i = \delta$ for $p + 1 \leq i \leq q - 1$ (for the neighbors of v_k)
- $\delta'_i = \delta_i$ for $q \leq i \leq t$.
- By induction hypothesis we can move w_1, \dots, w_t by $\delta'_1, \dots, \delta'_t$, respectively.
- We can complete the drawing by placing v_k , v_k is moved with $L(w_{p+1}), \dots, L(w_{q-1})$ by δ .

Algorithm Shift

Let v_1, \dots, v_n be a canonical ordering of G

for $i = 1$ **to** n **do**

└ $L(v_i) \leftarrow \{v_i\};$

$P(v_1) \leftarrow (0, 0); P(v_2) \leftarrow (2, 0); P(v_3) \leftarrow (1, 1);$

for $i = 4$ **to** n **do**

└ Let $w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2$ denote the boundary of $G_{i-1};$
and let w_p, \dots, w_q be the neighbors $v_i;$

└ **for** $\forall v \in \cup_{j=p+1}^{q-1} L(w_j)$ **do**

└ $x(v) \leftarrow x(v) + 1;$


└ **for** $\forall v \in \cup_{j=q}^t L(w_j)$ **do**

└ $x(v) \leftarrow x(v) + 2;$



└ $P(v_i) \leftarrow$ intersection of $+1$ and -1 edges from $P(w_p)$ and $P(w_q);$

└ $L(v_i) = \cup_{j=p+1}^{q-1} L(w_j) \cup \{v_i\};$



Linear Time Implementation of Shift Algorithm



Linear Time Implementation of Shift Algorithm



Linear Time Implementation of Shift Algorithm



- $x(v_k) = \frac{1}{2}(x(w_q) + x(w_p) + y(w_q) - y(w_p))$ (1)


- $y(v_k) = \frac{1}{2}(x(w_q) - x(w_p) + y(w_q) + y(w_p))$ (2)

- $x(v_k) - x(w_p) = \frac{1}{2}(x(w_q) - x(w_p) + y(w_q) - y(w_p))$ (3)


- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.

Linear Time Implementation of Shift Algorithm


- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- Calculate $\Delta_x(v_k)$ by eq. (3)
- Calculate $y(v_k)$ by eq. (2)




- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- Calculate $\Delta_x(v_k)$ by eq. (3)
- Calculate $y(v_k)$ by eq. (2)




- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- Calculate $\Delta_x(v_k)$ by eq. (3)
- Calculate $y(v_k)$ by eq. (2)



- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- Calculate $\Delta_x(v_k)$ by eq. (3)
- Calculate $y(v_k)$ by eq. (2)



- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- Calculate $\Delta_x(v_k)$ by eq. (3)
- Calculate $y(v_k)$ by eq. (2)



- If we know the y -coordinates of w_p and w_q and the difference $x(w_p) - x(w_q)$, we can compute the relative distance of v_k and w_p .
- In the binary tree which we construct we keep the relative x -distance of each node from its parent.
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- Calculate $\Delta_x(v_k)$ by eq. (3)
- Calculate $y(v_k)$ by eq. (2)

- $\Delta_x(w_q) = \Delta_x(w_p, w_q) - \Delta_x(v_k)$
- $\Delta_x(w_{p+1}) = \Delta_x(w_{p+1}) - \Delta_x(v_k)$

