

## Exercise Sheet 2

Discussion: 23. November 2017

### Exercise 1: Outerplanar and Series-Parallel Graphs ★

A graph  $G$  is called *outerplanar* if it has a planar drawing where all vertices lie on the boundary of the outer face. Prove the following lemma.

**Lemma 1** *Every biconnected outerplanar graph is series-parallel.*

### Exercise 2: Visibility Representation ★




Figure 1: A visibility representation (b) of the graph  $G$  (a).

In a *visibility representation* of a graph  $G = (V, E)$  the vertices are represented by horizontal segments (*vertex-segments*). We say that two vertices  $u$  and  $v$  *see* each other, if their vertex-segments can be connected by a vertical rectangle of non-zero width that does not cross any other vertex-segment. Thus, in a visibility representation of  $G$ , two vertices  $u, v$  see each other if and only if  $(u, v) \in E$ ; see Fig. 1. Prove the following lemma.

**Lemma 2** *Every series-parallel graph has a visibility representation.*

**Exercise 3: 2**

★

Let  $G = (V, E)$  be a triconnected plane graph with a vertex  $v_1$  on the outer face. Further, let  $\pi = (V_1, \dots, V_K)$  be an ordered partition of  $V$ , that is,  $V_1 \cup \dots \cup V_K = V$  and  $V_i \cap V_j = \emptyset$  for  $i \neq j$ . We define  $G_k$  to be the subgraph of  $G$  induced by  $V_1 \cup \dots \cup V_k$  and denote by  $C_k$  the outer face of  $G_k$ .

The sequence  $\pi$  is a *canonical ordering* of  $G$ , if

- $V_1$  consists of  $\{v_1, v_2\}$ , where  $v_2$  lies on the outer face and  $(v_1, v_2) \in E$ .
- $V_K = \{v_n\}$  is a singleton, where  $v_n$  lies on the outer face,  $(v_1, v_n) \in E$ , and  $v_n \neq v_2$ .
- Each  $C_k$  ( $k > 1$ ) is a cycle containing  $\{v_1, v_2\}$ .
- Each  $G_k$  is biconnected and internally triconnected, that is, removing two interior vertices of  $G_k$  does not disconnect it.
- For each  $k$  with  $2 \leq k \leq K - 1$ , one of the following conditions holds:
  1.  $V_k = \{z\}$ , where  $z$  belongs to  $C_k$  and has at least one neighbor in  $G - G_k$ .
  2.  $V_k = \{z_1, \dots, z_\ell\}$  is a chain, where each  $z_i$  has at least one neighbor in  $G - G_k$  and where  $z_1$  and  $z_\ell$  each have one neighbor on  $C_{k-1}$ , and these are the only two neighbors of  $V_k$  in  $G_{k-1}$ .

Prove the following lemma.

**Lemma 3** *Every triconnected planar graph admits a canonical ordering.*

**Hint:** Use reverse induction. For the induction step, consider the two cases that  $G_k$  is triconnected and  $G_k$  is not triconnected.

### Exercise 4: Barycentric Coordinates


★

Let  $\Delta_{a,b,c}$  be a triangle on the plane on vertices  $a$ ,  $b$  and  $c$ . For each point  $x$  laying inside triangle  $\Delta_{a,b,c}$  there exists a triple  $(x_a, x_b, x_c)$  such that  $x_a \cdot a + x_b \cdot b + x_c \cdot c = x$  and  $x_a + x_b + x_c = 1$ . The triple  $(x_a, x_b, x_c)$  is called *barycentric coordinates* of  $x$  with respect to  $\Delta_{a,b,c}$ .

Prove that:

- (a) If  $A(\Delta)$  denotes the area of the triangle  $A$ , then

$$x_a = \frac{A(\Delta_{b,c,x})}{A(\Delta_{a,b,c})}, \quad x_b = \frac{A(\Delta_{a,c,x})}{A(\Delta_{a,b,c})}, \quad x_c = \frac{A(\Delta_{a,b,x})}{A(\Delta_{a,b,c})}$$



- (b) Equations  $x_a = 0$ ,  $x_b = 0$ ,  $x_c = 0$  represent lines through  $bc$ ,  $ab$  and  $ac$ , respectively.
- (c) Let  $(x_a, x_b, x_c)$  be barycentric coordinates of point  $x$  in triangle  $\Delta_{abc}$ . The set of points  $\{(x_a, x'_b, x'_c) : x'_b, x'_c \in \mathbb{R}\}$  represents a line parallel to edge  $bc$  passing through point  $x$ . Similarly, sets of points  $\{(x'_a, x_b, x'_c) : x'_a, x'_c \in \mathbb{R}\}$ ,  $\{(x'_a, x'_b, x_c) : x'_a, x'_b \in \mathbb{R}\}$  represent lines parallel to edges  $ac$ ,  $ab$ , respectively, passing through point  $x$ .